



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

SOLUTIONS OF PROBLEMS IN NUMBER TWO.

SOLUTIONS of problems in No. 2 have been received as follows :

From R. J. Adcock, 299, 307, 308; Prof. L. G. Barbour, 296; Prof. W. P. Casey, 296, 299, 300, 301, 302; Alex. S. Christie, 301, 302; Prof. E. J. Edmunds, 296; George Eastwood, 298, 300; Dr. Wm. Hillhouse, 296, 300; W. E. Heal, 300, 302, 303, 305, 306; Prof. W. W. Johnson, 306; Chas. H. Kummell, 303, 308; Prof. J. H. Kershner, 296, 298, 299, 300, 301, 303; Prof. D. J. Mc Adam, 296; P. Richardson, 296, 300; T. P. Stowell, 298; Prof. E. B. Seitz, 296, 300, 307; Prof. J. Scheffer, 296, 298, 299, 300, 301, 303, 304; E. P. Thompson, 296.

296. "Given  $x^6 + y^6 = \frac{1}{2}\sqrt{2}$ ,  $x^4 + y^4 = 1$ ; to find  $x$  and  $y$  by quadratics."

SOLUTION BY PROF. L. G. BARBOUR, RICHMOND, KY.

Let  $m+n = x^2$ ,  $m-n = y^2$ ; then

$$m^3 + 3m^2n + 3mn^2 + n^3 + m^3 - 3m^2n + 3mn^2 - n^3 = x^6 + y^6 = \frac{1}{2}\sqrt{2}.$$

$$\therefore m^3 + 3mn^2 = \frac{1}{2} \cdot \frac{1}{2}\sqrt{2} = \frac{1}{4}\sqrt{2}. \quad (1)$$

Also  $m^2 + 2mn + n^2 + m^2 - 2mn + n^2 = x^4 + y^4 = 1$ ;  $\therefore m^2 + n^2 = \frac{1}{2}$ .

$\therefore n^2 = \frac{1}{2} - m^2 =$  by (1),  $(\frac{1}{4}\sqrt{2} - m^3) \div 3m$ , or  $\frac{3}{2}m - 3m^2 = \frac{1}{4}\sqrt{2} - m^3$ .

$$\therefore 4m^3 - 3m = -\frac{1}{2}\sqrt{2}, \therefore 4m^4 - 3m^2 = -\frac{1}{2}\sqrt{2} \cdot m,$$

$$\text{or} \quad 4m^4 - m^2 + \frac{1}{16} = 2m^2 - \frac{1}{2}\sqrt{2} \cdot m + \frac{1}{16};$$

$$\therefore 2m^2 - \frac{1}{2} = m\sqrt{2} - \frac{1}{4}; \therefore 2m = \sqrt{2}, \text{ or } m = \frac{1}{2}\sqrt{2}.$$

Therefore  $n^2 = 0$ , and  $x^4 = m^2 = \frac{1}{2}$ ;  $\therefore x = \frac{1}{2}\sqrt{2} = y$ .

[This problem was also solved as above, with a slight difference in notation, by Professors Seitz and Mc Adam.]

297. No solution received.

298. "On the diameter  $AB$  produced, of a given circle, is a given point  $P$ . A chord  $CD$  is parallel to the diameter  $AB$ . Find the position of  $CD$ , by a geometrical construction, such that the angle  $CPD$  shall be the greatest possible."

SOLUTION BY PROF. J. H. KERSHNER, MERCERSBURG, PA.

Represent the point  $D$  by  $(x, y)$ , the origin being at the centre. The tan of  $CPD = \tan(CPB - DPB) = (\tan CPB - \tan DPB) \div (1 + \tan CPB \times \tan DPB)$

$$= \left( \frac{y}{a-x} - \frac{y}{a+x} \right) \div \left( 1 - \frac{y^2}{a^2-x^2} \right)$$


$$= \left( \frac{2xy}{a^2-x^2+y^2} \right) = \text{a maximum,}$$

where  $a = PO$ .

Substituting  $\sqrt{(r^2-x^2)}$  for  $y$ , we find by the Diff. Calculus, the expression has a maximum for

$$x = \frac{r\sqrt{(a^2-r^2)}}{a\sqrt{2}}.$$

To find  $x$  by geometrical construction, erect the perpendicular  $OF=OP$ , join  $PF$  and  $PH$ , lay off  $FK=r$ , draw  $KL$ , parallel to  $OF$ , meeting  $PH$  in  $L$ , and we shall have  $HL=x$  = the abscissa of  $C$  or  $D$ .



299. "Prove that the number of terms in an *expanded* polynomial of  $t$  terms is equal to the number of combinations of  $(n+t-1)$  quantities taken  $t-1$  at a time, where  $n$  is the exponent of the polynomial."

SOLUTION BY PROF. W. P. CASEY, SAN FRANCISCO, CAL.

The number of terms in the expansion of any polynomial, the exponent  $n$ , being a positive integer, is

$$\frac{t(t+1)(t+2)\dots(t+n-1)}{n!},$$

where  $t$  is the number of terms and  $n$  the exponent; which is evidently = to the number of combinations of  $(n+t-1)$  quantities taken  $t-1$  at a time.

SOLUTION BY R. J. ADCOCK, ROSEVILLE, ILL.

The expansion of  $(a+b)^n$  has  $n+1$  terms; that of  $(a+b+c)^n$  =  $a^n + na^{n-1}(b+c) + \frac{1}{2}n(n-1)a^{n-2}(b+c)^2 + \dots + na(b+c)^{n-1} + (b+c)^n$ , which has the sum of  $n+1$  terms of the series  $1+2+3+4+\dots+(n+1)$  =  $\frac{1}{2}(n+2)(n+1)$  terms. The expansion of  $(a+b+c+d)^n$  has the sum of  $n+1$  terms of the series  $1+3+6+\dots + \frac{(n+1)(n+2)}{1.2} = \frac{(n+3)(n+2)(n+1)}{1.2.3}$

by differences. In like manner, the expansion of  $(a+b+c+d+e)^n$  has  $\frac{(n+4)(n+3)(n+2)(n+1)}{1.2.3.4}$  terms. Hence  $\frac{(n+t-1)(n+t-2)\dots(n+1)}{1.2.3\dots(t-1)}$

which expresses the proposition that was to be proved.

300. "In a plane trian.  $ABC$  are given, the perimeter  $2s$  and the radii of the inscribed and circumscribed circles,  $r$  and  $R$ , to determine the triangle."

SOLUTION BY P. RICARDSON, BROOKLYN, N. Y.

Let the sides opposite the angles  $A, B, C$ , be  $a, b, c$ , then we have given

$$a + b + c = 2s. \quad (1)$$

Double the area is  $ab \sin C = 2rs$ ,  $\sin C = c \div 2R$ ;  $\therefore$  by substitution we get

$$abc = 4Rrs. \quad (2)$$

The area is  $\sqrt{s(s-a)(s-b)(s-c)} = rs$ , or, by squaring and performing operations indicated in left hand member, we get

$$s^4 - (a+b+c)s^3 + (ab+ac+bc)s^2 - abcs = r^2s^2,$$

and by substituting from (1) and (2) we get

$$[-s^4 + (ab+ac+bc)s^2 - 4Rrs^2] = r^2s^2,$$

whence

$$ab+ac+bc = r^2+s^2+4Rr. \quad (3)$$

Equations (1), (2) and (3) show that  $a, b, c$ , must be the roots of the cubic equation  $x^3 - 2sx^2 + (r^2 + s^2 + 4Rr)x - 4Rrs = 0$ .

301. "Determine the angles of a plane triangle from the following relations.

$$(I) \begin{cases} \tan^2 A + \tan^2 B + \tan^2 C = m, \\ \tan^4 A + \tan^4 B + \tan^4 C = n. \end{cases} \quad (1)$$

$$(II) \begin{cases} \tan^{\frac{1}{2}} A + \tan^{\frac{1}{2}} B + \tan^{\frac{1}{2}} C = m, \\ \tan^{\frac{1}{4}} A + \tan^{\frac{1}{4}} B + \tan^{\frac{1}{4}} C = n. \end{cases} \quad (1)$$

$$(III) \begin{cases} \tan \frac{1}{2} A + \tan \frac{1}{2} B + \tan \frac{1}{2} C = m, \\ \tan \frac{1}{2} A \times \tan \frac{1}{2} B \times \tan \frac{1}{2} C = n. \end{cases} \quad (1)$$

SOLUTION BY ALEX. S. CHRISTIE, U. S. C. SURV., WASH., D. C.

Let the sought tangents in each case be the roots  $a, b, c$  of the cubic

$$x^3 + px^2 + qx + r = 0.$$

Then by well known forms in the theory of equations and trigonometry (say Faa' de Bruno's Forms, Binaires, p. 27, and Chauvenet's Trig., p. 62) we have for determining  $p, q, r$ ,

$$(I) \begin{cases} m = a^2 + b^2 + c^2 = p^2 - 2q \\ n = a^4 + b^4 + c^4 = p^4 - 4p^2q + 4pr + 2q^2 \\ 0 = p - r \end{cases} \quad (1)$$

$$(II) \quad (1) \text{ and } (2) \text{ with } 0 = 1 - q$$

$$(III) \quad \begin{cases} m = -p \\ n = -r \\ 0 = 1 - q \end{cases}$$

Whence, for the three cases in order





304. "Find the limit of error in the following method of trisecting an arc approximately: Trisect the chord of the arc, also the diameter parallel to it, and the corresponding semicircumference. Join the corresponding points of trisection of the semicircumference and the diameter, producing the lines till they intersect. From the point thus found pass lines through the points of trisection of the chord until they meet the arc."


SOLUTION BY PROF. J. SCHEFFER, MERCERSBURG, PA.

Let  $E, E'$  represent the points of trisection of the diameter  $MN$ ;  $C, C'$ , those of the chord  $AB$ , and  $F, F'$ , those of the semicircle.

Draw the perpendicular  $PD$ , and join the centre  $O$  with  $F$  and  $G$ . Denote the angle  $AOD$  by  $\alpha$ , the angle  $GOH$  by  $\varphi$ , and the radius by  $r$ .

We now easily find  $OE = \frac{1}{3}r$   $CD = \frac{1}{3}\sin \alpha$ .

In the triangle  $EOF$ , we have  $EF^2 = EO^2 + OF^2 - 2EO \cdot OF \cos 60^\circ$ ;  $\therefore EF = \frac{1}{3}r\sqrt{7}$ ;  $\cos FEO = (FE^2 + EO^2 - OF^2) \div (2EF \cdot EO) = -\frac{1}{14}\sqrt{7}$ ;  $\therefore OEP = \frac{1}{14}\sqrt{7}$ , and  $\sin OEP = \frac{3}{14}\sqrt{21}$ ; therefore  $\tan OEP = 3\sqrt{3}$ . But  $PO = \frac{1}{3}r \times \tan OEP = r\sqrt{3}$ . Since  $OD = r \cos \alpha$ , we have  $PD = r(\cos \alpha + \sqrt{3})$ .



We have  $\tan GPO = CD \div PD = \frac{1}{3}[\sin \alpha \div (\cos \alpha + \sqrt{3})]$ ; therefore  $\sin GPO = \frac{\sin \alpha}{\sqrt{[\sin^2 \alpha + 9(\cos \alpha + \sqrt{3})^2]}}$ ;  $\cos GPO = \frac{3(\cos \alpha + \sqrt{3})}{\sqrt{[\sin^2 \alpha + 9(\cos \alpha + \sqrt{3})^2]}}$

In the triangle  $GPO$ , we have  $OG : OP = \sin GPO : \sin PGO$ ;  $\therefore \sin PGO = \frac{\sqrt{3} \cdot \sin \alpha}{\sqrt{[\sin^2 \alpha + 9(\cos \alpha + \sqrt{3})^2]}}$ ;  $\cos PGO = \frac{\sqrt{[9(\cos \alpha + \sqrt{3})^2 - 2\sin^2 \alpha]}}{\sqrt{[\sin^2 \alpha + 9(\cos \alpha + \sqrt{3})^2]}}$

But  $\sin \varphi = \sin (GPO + PGO)$ , therefore

$$\sin \varphi = \frac{\sin \alpha \sqrt{[9(\cos \alpha + \sqrt{3})^2 - 2\sin^2 \alpha]} + 3\sqrt{3} \cdot \sin \alpha (\cos \alpha + \sqrt{3})}{\sin^2 \alpha + 9(\cos \alpha + \sqrt{3})^2}$$

With regard to the trisection of an arc this formula gives a correct result only for an arc of  $180^\circ$ . For other arcs, I find, if  $\alpha = 45^\circ$ ,  $\varphi = 15^\circ 6'$ ; if  $\alpha = 30^\circ$ ,  $\varphi = 10^\circ 2'$ , &c.

305. "Find the equation of the curve, rectangular co-ordinates  $(x, y)$  in which the length  $s = \frac{ny}{x} = \int_0^x dx \sqrt{[1 + (\frac{dy}{dx})^2]}$ ,  $n$  being a constant."

SOLUTION BY W. E. HEAL, WHEELING, INDIANA.

Let the indefinite integral

$$\int dx \left[ 1 + \left( \frac{dy}{dx} \right)^2 \right]^{\frac{1}{2}} = \varphi(x), \text{ then } \int_0^x dx \left[ 1 + \left( \frac{dy}{dx} \right)^2 \right]^{\frac{1}{2}} = \varphi(x) - \varphi(0) = \frac{ny}{x}$$

Differentiating the above eq'n, (1), remembering that  $\varphi(0)$  is constant,

$$dx \sqrt{1 + \left( \frac{dy}{dx} \right)^2} = \frac{n(xdy - ydx)}{x^2}. \quad (2)$$

Put  $dy \div dx = p$ , and (2) becomes  $x^2 \sqrt{1 + p^2} = npy - ny$ . (3)

Differentiating (3) we have

$$\frac{xpd p}{\sqrt{1 + p^2}} + 2dx \sqrt{1 + p^2} = ndp. \quad (4)$$

Divide by  $2(1 + p^2)^{\frac{1}{2}}$ ;  $dx(1 + p^2)^{\frac{1}{2}} + xpd p \div (1 + p^2)^{\frac{1}{2}} = ndp \div 2(1 + p^2)^{\frac{1}{2}}$ . (5)

Integrating (5) and determining the constant  $C = 0$ , we have

$$x(1 + p^2)^{\frac{1}{2}} = \frac{n}{2} \int \frac{dp}{(1 + p^2)^{\frac{1}{2}}}, \quad (6). \quad \text{Let } p = \frac{2^{\frac{1}{2}}v(1 - \frac{1}{2}v^2)^{\frac{1}{2}}}{1 - v^2}, \text{ then (6) becomes}$$

$$\begin{aligned} \frac{x}{(1 - v^2)^{\frac{1}{2}}} &= \frac{n}{2^{\frac{1}{2}}} \int \frac{dv}{\sqrt{[(1 - v^2)^3(1 - \frac{1}{2}v^2)]}} = 2^{\frac{1}{2}}n \left[ \frac{v(1 - \frac{1}{2}v^2)^{\frac{1}{2}}}{(1 - v^2)^{\frac{1}{2}}} \right. \\ &+ \frac{1}{2} \int \frac{dv}{(1 - v^2)^{\frac{1}{2}}(1 - \frac{1}{2}v^2)^{\frac{1}{2}}} - \left. \int \frac{(1 - \frac{1}{2}v^2)^{\frac{1}{2}}dv}{(1 - v^2)^{\frac{1}{2}}} \right] = 2^{\frac{1}{2}}n \left[ \frac{v(1 - \frac{1}{2}v^2)^{\frac{1}{2}}}{(1 - v^2)^{\frac{1}{2}}} + \frac{1}{2}F(\frac{1}{2}2^{\frac{1}{2}}, v) \right. \\ &- \left. E(\frac{1}{2}2^{\frac{1}{2}}, v) \right] \quad (7), \text{ where } F \text{ and } E \text{ are elliptic funct's of the 1st and 2nd ord's.} \end{aligned}$$

But  $v = \frac{[(1 + p^2)^{\frac{1}{2}} - 1]^{\frac{1}{2}}}{(1 + p^2)^{\frac{1}{2}}}$ , and from (3)  $p = \frac{n^2y + x[n^2(x^2 + y^2) - x^4]}{x(n^2 - x^2)}$ .

Substituting these values in (7) we have the equation of the curve.

[The solution of 306, by Prof. Johnson, of 307, by Prof. Seitz, and of 308, by Mr. Adcock, will be published in No. 4.]

### PROBLEMS.

309. *By Prof. Beman.*—In a given circle, find the vertices of the inscribed square, pentagon, octagon, and decagon by using the dividers alone.

310. *By Prof. Edmunds.*—Required the locus of vertices of a right angled spherical triangle whose legs pass through two fixed points given on the surface of the sphere.

311. *By Prof. Casey.*—A uniform circular plate is placed with its centre upon a prop, to find at what points on its circumference three given weights  $p, q, r$  must be attached that it may remain at rest in a horizontal position.

312. *By Prof. Scheffer.*—To find the area of the loop of the curve  $y^3 + x^2y - axy + bx^2 = 0$ .